

ON LINEAR DEPENDENCE OF ITERATES

J. MATKOWSKI and W. ZHANG

Received September 9, 1997 and, in revised form, February 1, 2000

Abstract. A functional equation related to a problem of linear dependence of iterates is considered.

1. Introduction

The polynomial-like iterative functional equation

$$\lambda_0 f^0(x) + \lambda_1 f^1(x) + \dots + \lambda_n f^n(x) = F(x), \quad x \in X,$$

where X stands for a real or complex linear space and f^k denotes the k -th iterate of the unknown function $f : X \rightarrow X$, i.e., $f^0(x) = x$ for $x \in X$ and $f^{k+1} = f \circ f^k$ (here " \circ " denotes the composition of functions) is discussed extensively, cf. [1]–[11]. An important special case of this equation is

$$f^n(x) = a_{n-1}f^{n-1}(x) + a_{n-2}f^{n-2}(x) + \dots + a_0x, \quad x \in X, \quad (1)$$

where a_0, \dots, a_{n-1} are real or complex numbers. This functional equation can be interpreted as linear dependence of iterates of f . In 1974 Nabeya

1991 *Mathematics Subject Classification.* 39B12.

Key words and phrases. Iterate functional equation, characteristic polynomial.

[8] discussed (1) for $n = 2$ and $\mathbf{X} = \mathbf{R}$ in detail by considering its characteristic equation. However Nabeya's idea appears to be difficult to apply in solving equation (1) for $n \geq 3$. During the 26th International Symposium on Functional Equations held in Spain in 1988 the first author presented the result [6] that *the solutions of (1) for $n = k$ are solutions of (1) for $n = m$, $m \geq k$, if the characteristic polynomial of the lower order equation exactly divides that of the higher order one*. This statement establishes a useful relation in the class of iterative equations of type (1), but until now the proof was not published. In this paper an elementary proof is presented. Furthermore, based on this result some conclusions how the solutions to be ruled by the roots of the relevant characteristic polynomials are given.

2. Characteristic equations

Following Euler's idea for differential equations, we formally consider a linear solution

$$f(x) = rx, \quad x \in \mathbf{X},$$

of the equation (1) where $r \in \mathbf{C}$ is indeterminate. From (1) we have

$$r^n - a_{n-1}r^{n-1} - \dots - a_1r - a_0 = 0. \quad (2)$$

Here (2) is called the *characteristic equation* of equation (1), its roots are called the *characteristic roots*, and the left-hand side of (2), denoted by $P_n(r)$, is called the *characteristic polynomial* of equation (1). By the well known relations between roots and coefficients of polynomials equation (1) is equivalent to

$$f^n(x) - \left(\sum_{i=1}^n r_i\right)f^{n-1}(x) + \left(\sum_{i < j} r_i r_j\right)f^{n-2}(x) + \dots + (-1)^n r_1 r_2 \dots r_n x = 0 \quad (3)$$

for $x \in \mathbf{X}$, where r_1, r_2, \dots, r_n are n complex roots of the polynomial P_n . Let $F_n(r_1, r_2, \dots, r_n)f$ denote the function of the left-hand side of (3) and call it n -form of (3). The n -form is uniquely determined by given $r_1, r_2, \dots, r_n \in \mathbf{C}$.

Lemma 1. For fixed $r_1, r_2, \dots, r_{n+1} \in \mathbf{C}$, if $F_n(r_1, r_2, \dots, r_n)f = 0$ then

$$F_{n+1}(r_1, \dots, r_n, r_{n+1})f = 0.$$

Proof. Since $F_n(r_1, r_2, \dots, r_n)f = 0$, i.e., f satisfies equation (3), we have

$$\begin{aligned} f^{n+1}(x) = f^n(f(x)) &= \left(\sum_{i=1}^n r_i\right)f^n(x) - \left(\sum_{i < j} r_i r_j\right)f^{n-1}(x) + \dots \\ &\quad + (-1)^{n+1} r_1 r_2 \dots r_n f(x), \quad x \in \mathbf{X}. \end{aligned}$$

Thus, for all $x \in X$, the $(n+1)$ -form satisfies

$$\begin{aligned} & F_{n+1}(r_1, \dots, r_n, r_{n+1})f(x) \\ &= f^{n+1}(x) - \left(\sum_{i=1}^{n+1} r_i\right)f^n(x) + \left(\sum_{i < j}^{n+1} r_i r_j\right)f^{n-1}(x) + \dots + (-1)^{n+1} r_1 r_2 \dots r_{n+1} x \\ &= \left(\sum_{i=1}^n r_i - \sum_{i=1}^{n+1} r_i\right)f^n(x) - \left(\sum_{i < j}^n r_i r_j - \sum_{i < j}^{n+1} r_i r_j\right)f^{n-1}(x) \\ &+ \dots + (-1)^{n+1} r_1 r_2 \dots r_{n+1} x \\ &= -r_{n+1} f^n(x) + r_{n+1} \left(\sum_{i=1}^n r_i\right)f^{n-1}(x) - r_{n+1} \left(\sum_{i < j}^n r_i r_j\right)f^{n-2}(x) \\ &+ \dots + (-1)^{n+1} r_1 r_2 \dots r_{n+1} x = -r_{n+1} F_n(r_1, r_2, \dots, r_n)f(x) = 0. \end{aligned}$$

□

Now we can prove the result presented in [6].

Theorem 1. Suppose that

$$Q(r) = r^k - b_{k-1}r^{k-1} - \dots - b_1r - b_0,$$

$$P(r) = r^n - a_{n-1}r^{n-1} - \dots - a_1r - a_0,$$

are polynomials, where $r \in \mathbb{C}$, $k \leq n$, and that $Q|P$, i.e., P is exactly divided by Q . If a function $f: X \rightarrow X$ satisfies the functional equation

$$f^k(x) = b_{k-1}f^{k-1}(x) + b_{k-2}f^{k-2}(x) + \dots + b_0x, \quad x \in X, \quad (4)$$

then f satisfies functional equation (1), i.e.,

$$f^n(x) = a_{n-1}f^{n-1}(x) + a_{n-2}f^{n-2}(x) + \dots + a_0x, \quad x \in X.$$

Proof. Let r_1, r_2, \dots, r_n be complex roots of P . Since $Q|P$ we may assume without any loss of generality that r_1, \dots, r_k , $k \leq n$, are roots of Q . From the definition of F_k and (4) we have

$$F_k(r_1, r_2, \dots, r_k)f = 0.$$

By Lemma 1, the function f also satisfies

$$F_{k+1}(r_1, \dots, r_k, r_{k+1})f = 0.$$

Thus, by induction, we can prove easily that

$$F_n(r_1, r_2, \dots, r_n)f = 0,$$

that is, f satisfies equation (1). □

Remark 1. Equation (1) of order n has a solution which does not satisfy the equation (4) of order k if $Q|P$ but $Q \neq P$.

In fact, if all roots r_1, r_2, \dots, r_n of P are real and only r_1, r_2, \dots, r_k , $k < n$, are roots of Q , then $f(x) = r_i x$, $x \in X$, $i = k+1, \dots, n$, satisfies (1) but is not a solution of (4).

Remark 2. Let $X = \mathbb{R}$ and suppose that the coefficients in equation (1) are real. If r_0 is a complex root of the characteristic polynomial P_n with imaginary part $\Im r_0 \neq 0$, then all solutions of the real 2-order iterative equation

$$f^2(x) = 2\Re r_0 f(x) - |r_0|^2 x,$$

where $\Re r_0$ denotes the real part of r_0 and $|r_0|$ denotes the modulus of r_0 , satisfy equation (1).

This assertion is a consequence of Theorem 1 and the fact that the conjugacy \bar{r}_0 of r_0 is also a root of P_n .

3. Iterations of solutions

For convenience, let $F_{n-1}(r_1, \dots, \check{r}_k, \dots, r_n)f$ represent the $(n-1)$ -form of (3) determined by $n-1$ characteristic roots $r_1, \dots, r_{k-1}, r_{k+1}, \dots, r_n$.

Theorem 2. Suppose that the characteristic polynomial P_n in (2) has n pairwise different roots r_1, \dots, r_n and that $f : X \rightarrow X$ is a solution of functional equation (1). Then for any integer $m \geq 0$,

$$f^{n+m} = \frac{A_{11}}{\Delta} r_1^{m+1} g_1 + \frac{A_{21}}{\Delta} r_2^{m+1} g_2 + \dots + \frac{A_{n1}}{\Delta} r_n^{m+1} g_n, \quad (5)$$

where

$$g_k := F_{n-1}(r_1, \dots, \check{r}_k, \dots, r_n)f, \quad k = 1, 2, \dots, n,$$

and Δ and A_{k1} , $k = 1, 2, \dots, n$, denote respectively the determinant and algebraic adjuncts of the matrix

$$A = \begin{pmatrix} 1 - \sum_{i \neq 1} r_i & \sum_{i < j, \neq 1} r_i r_j & \dots & (-1)^{n-1} r_2 r_3 \dots r_n \\ 1 - \sum_{i \neq 2} r_i & \sum_{i < j, \neq 2} r_i r_j & \dots & (-1)^{n-1} r_1 r_3 \dots r_n \\ \dots & \dots & \dots & \dots \\ 1 - \sum_{i \neq n} r_i & \sum_{i < j, \neq n} r_i r_j & \dots & (-1)^{n-1} r_1 r_2 \dots r_{n-1} \end{pmatrix}. \quad (6)$$

Here $\sum_{i \neq 1}$ and its like denote the summations with respect to the indexes from 1 to n with some shown restriction.

Proof. Write equation (3) in the equivalent form

$$\begin{aligned} f^n - \left(\sum_{i \neq n} r_i \right) f^{n-1} + \left(\sum_{i < j, i \neq n} r_i r_j \right) f^{n-2} + \dots + (-1)^{n-1} r_1 r_2 \dots r_{n-1} f \\ = r_n f^{n-1} - r_n \left(\sum_{i \neq n} r_i \right) f^{n-2} + \dots + (-1)^{n-1} r_1 r_2 \dots r_n f^0. \end{aligned}$$

By the definition of g_k , with $k = n$, we have

$$g_n \circ f = r_n g_n.$$

Thus, for every non-negative integer m ,

$$g_n \circ f^{m+1} = r_n^{m+1} g_n,$$

that is,

$$f^{n+m} - \left(\sum_{i \neq n} r_i \right) f^{n+m-1} + \dots + (-1)^{n-1} r_1 r_2 \dots r_{n-1} f^{m+1} = r_n^{m+1} g_n,$$

is a linear equation for $f^{n+m}, f^{n+m-1}, \dots, f^{m+1}$. Similarly, for each fixed k , $k = 1, 2, \dots, n-1$, we get another linear equation. Thus we obtain a system of n linear equations, expressed by

$$AF = G,$$

where A is a matrix defined by (6), F and G are transposes of the vectors $(f^{n+m}, f^{n+m-1}, \dots, f^{m+1})$ and $(r_1^{m+1} g_1, r_2^{m+1} g_2, \dots, r_n^{m+1} g_n)$, respectively. Applying repeatedly elementary linear transformations on the rows of A we obtain

$$\Delta = \det A = \prod_{i < j, i=1}^n (r_j - r_i) \neq 0,$$

i.e., A is invertible. Now formula (5) is a direct consequence of Cramer's rule. \square

Corollary 1. Suppose that the polynomial P_n in (2) has n pairwise different roots r_1, \dots, r_n and that $f: X \rightarrow X$ is a solution of a k -order equation of the form (1) whose characteristic polynomial Q_k exactly divides P_n . Then f^{n+m} is a sum of the suitable k terms which appear in (5).

Proof. Since $Q_k | P_n$, we may assume without any loss of generality that the first k numbers r_1, r_2, \dots, r_k are the k roots of Q_k . Thus the function f satisfies the equation $F_k(r_1, r_2, \dots, r_k)f = 0$. By Theorem 1,

$$F_{n-1}(r_1, \dots, r_k, \dots, \bar{r}_i, \dots, r_n)f = 0, \quad i = k+1, \dots, n,$$

that is, according to the notations in Theorem 2, $g_i = 0$, $i = k + 1, \dots, n$. By Theorem 2,

$$f^{n+m} = \frac{A_{11}}{\Delta} r_1^{m+1} g_1 + \frac{A_{21}}{\Delta} r_2^{m+1} g_2 + \dots + \frac{A_{k1}}{\Delta} r_k^{m+1} g_k, \quad m \geq 0,$$

which completes the proof. \square

Remark 3. It is easy to verify that a solution $f : X \rightarrow X$ of (1) is one-to-one if $a_0 \neq 0$; if moreover $X = \mathbf{R}$ and f is continuous then it is strictly monotone and onto. If $a_0 \neq 0$ then, by (2), the characteristic polynomial of equation (1) has no zero root.

Obviously, if $a_0 \neq 0$ and f is onto then equations (1) and (3) are equivalent, respectively, to

$$f^{-n}(x) = -\frac{a_1}{a_0} f^{-(n-1)}(x) + \dots - \frac{a_{n-1}}{a_0} f^{-1}(x) + \frac{1}{a_0} x, \quad x \in X, \quad (7)$$

and

$$f^{-n} - \left(\sum_{i=1}^n s_i \right) f^{-(n-1)} + \left(\sum_{i < j} s_i s_j \right) f^{-(n-2)} + \dots + (-1)^n s_1 s_2 \dots s_n f^0 = 0, \quad (8)$$

where f^{-k} denotes the k -th iterate of f^{-1} and $s_i = r_i^{-1}$, $i = 1, 2, \dots, n$. In fact, in this case f is invertible, maps X onto itself and satisfies (1). Usually (7) and (8) are called the *dual equations* of (1) and (3), respectively. The following result is the dual counterpart of Theorem 2.

Theorem 3. Suppose that the hypotheses of Theorem 2 hold. If f is onto and $a_0 \neq 0$ in (1) then, for any integer $m \geq 0$,

$$f^{-(n+m)} = \frac{\tilde{A}_{11}}{\tilde{\Delta}} s_1^{m+1} \tilde{g}_1 + \frac{\tilde{A}_{21}}{\tilde{\Delta}} s_2^{m+1} \tilde{g}_2 + \dots + \frac{\tilde{A}_{n1}}{\tilde{\Delta}} s_n^{m+1} \tilde{g}_n,$$

where \tilde{g}_i , $\tilde{\Delta}$ and \tilde{A}_{i1} , $i = 1, 2, \dots, n$, are just modified g_i , Δ and A_{i1} , $i = 1, 2, \dots, n$, defined in Theorem 2 where r_j is replaced by s_j , $j = 1, 2, \dots, n$, and f is replaced by f^{-1} .

4. Some properties of solutions

Assume that X is a normed space, $a_0 \neq 0$ and that the characteristic polynomial of equation (1) has n pairwise different roots r_1, r_2, \dots, r_n .

Corollary 2. *Let $f : X \rightarrow X$ be a continuous solution of equation (1).*

- 1° *If $|r_k| < 1$ for all $k = 1, \dots, n$, then f^k approaches 0 as $k \rightarrow +\infty$;*
- 2° *If f is onto and $|r_k| > 1$ for all $k = 1, \dots, n$, then f^k approaches 0 as $k \rightarrow -\infty$;*
- 3° *In both cases 0 is a unique fixed point of f .*

Proof. Letting $m \rightarrow +\infty$ in (5) gives 1°. Similarly 2° is a consequence of the formula in Theorem 3. To prove 3° assume that $f(x_0) = x_0$ for some $x_0 \neq 0$. From (3),

$$x_0 - \left(\sum_{i=1}^n r_i\right)x_0 + \left(\sum_{i < j} r_i r_j\right)x_0 + \dots + (-1)^n r_1 r_2 \dots r_n x_0 = 0,$$

that is, $\prod_{i=1}^n (1 - r_i) = 0$. Thus at least one of r_i , $i = 1, 2, \dots, n$, would be equal 1. This contradicts the hypotheses in 1° and 2°. Therefore f has no non-zero fixed point.

Now the relation $f(0) = 0$ is an obvious consequence of the continuity of the function f . \square

In the next result we assume that $X = \mathbb{R}$.

Corollary 3. *Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is a strictly increasing and continuous solution of equation (1). The following results are true.*

- 1° *If $-1 < r_1 < \dots < r_{n-1} < 1 < r_n$ or $r_1 < -1 < r_2 < \dots < r_n < 1$, and if $f(x) < x$ for all $x > 0$ and $f(x) > x$ for all $x < 0$, then f satisfies*

$$F_{n-1}(r_1, \dots, r_{n-1})f = 0 \quad \text{or} \quad F_{n-1}(r_2, \dots, r_n)f = 0.$$

- 2° *If $r_1 < \dots < r_{n-1} < -1 < r_n$ or $r_1 < 1 < r_2 < \dots < r_n$, and if $f(x) > x$ for all $x > 0$ and $f(x) < x$ for all $x < 0$, then f satisfies*

$$F_{n-1}(r_1^{-1}, \dots, r_{n-1}^{-1})f^{-1} = 0 \quad \text{or} \quad F_{n-1}(r_2^{-1}, \dots, r_n^{-1})f^{-1} = 0.$$

Proof. By similar arguments as in the last corollary it is easy to show that, in both cases, 0 is a unique fixed point of f in \mathbf{R} . To prove 1^0 assume that $-1 < r_1 < \dots < r_{n-1} < 1 < r_n$ and take arbitrary $x > 0$. Since f is increasing, we have

$$x > f(x) > f^2(x) > \dots > f^k(x) \rightarrow 0, \quad \text{as } k \rightarrow +\infty, \quad x > 0.$$

Similarly, for arbitrary $x < 0$, we have

$$x < f(x) < f^2(x) < \dots < f^k(x) \rightarrow 0, \quad \text{as } k \rightarrow +\infty, \quad x < 0.$$

By Theorem 2, g_n vanishes, i.e., $F_{n-1}(r_1, \dots, r_{n-1})f = 0$ because $|r_i|^k \rightarrow 0, i = 1, 2, \dots, n-1$, and $|r_n|^k$ does not as $k \rightarrow +\infty$. Similarly, applying Theorem 3, we can prove 2^0 . \square

Acknowledgement. The authors are grateful to the referees for their helpful comments.

References

- [1] Fort Jr., M.K., *Continuous solutions of a functional equation*, Ann. Polon. Math. **13** (1963), 205–211.
- [2] Jarczyk, W., *On an equation of linear iteration*, Aequationes Math. **51** (1996), 303–310.
- [3] Jarczyk, W., *Series of iterates summing up to the identity function*, Grazer Math. Ber. **334** (1997), 153–162.
- [4] Kuczma, M., *Functional Equations in a Single Variable*, Monografie Matematyczne **46**, Warszawa, 1968.
- [5] Kuczma, M., Choczewski, B. and Ger, R., *Iterative Functional Equations*, Encyclopedia Math. Appl. **32**, Cambridge Univ. Press, 1990.
- [6] Matkowski, J., *The 26th International Symposium on Functional Equations, Remark 35*, Aequationes Math. **37** (1989), 119–120.
- [7] Mukherjea, A. and Ratti, J.S., *On a functional equation involving iterates of a bijection on the unit interval*, Nonlinear Anal. **7** (1983), 899–908.
- [8] Nabeya, S., *On the functional equation $f(p + qx + rf(x)) = a + bx + cf(x)$* , Aequationes Math. **11** (1974), 199–211.
- [9] Tabor, J. and Tabor, J., *On a linear iterative equation*, Results Math. **27** (1995), 412–421.
- [10] Zhang, W., *Discussion on the differentiable solutions of the iterated equation $\sum_{i=1}^n \lambda_i f^i(x) = F(x)$* , Nonlinear Anal. **15** (1990), 387–398.
- [11] Zhang, W. and Baker, J.A., *Continuous solutions of a polynomial-like iterative equation with variable coefficients*, Waterloo Rep. Fac. **11** (1994), University of Waterloo, Ontario, Canada.

JANUSZ MATKOWSKI
INSTITUTE OF MATHEMATICS
PEDAGOGICAL UNIVERSITY
PL. SŁOWIAŃSKI 9
65-069 ZIELONA GÓRA, POLAND
AND
INSTITUTE OF MATHEMATICS
SILESIA UNIVERSITY
KATOWICE, POLAND

WEINIAN ZHANG
DEPARTMENT OF MATHEMATICS
SICHUAN UNION UNIVERSITY
EAST AREA, CHENGDU 610064
P.R. CHINA